

Annex 8

REPORT OF THE BILLFISH WORKING GROUP WORKSHOP

International Scientific Committee for Tuna and Tuna-like Species
in the North Pacific Ocean

12-13 July 2010
Victoria, British Columbia, Canada

1.0 INTRODUCTION

An intercessional workshop of the Billfish Working Group (BILLWG) of the International Scientific Committee for Tuna and Tuna-like Species in the North Pacific Ocean (ISC) was convened in Victoria, British Columbia, Canada from 12-13 July 2010. The goals of this workshop were to 1) finalize advice on potential billfish biological reference points, 2) review the spatial extent and disposition of fisheries catching North Pacific striped marlin within areas (Western Central Pacific Ocean (WCPO) and Eastern Pacific Ocean (EPO)) delineated at the April 2010 BILLWG Workshop, 3) review and modify (if necessary) current conservation advice, and 4) review and discuss the World Blue Marlin Symposium proposal.

Gerard DiNardo, Chairman of the BILLWG, welcomed participants from Japan, Korea, and the United States of America (USA) (Attachment 1). Gerard DiNardo, Chair of the ISC BILLWG, provided the welcoming remarks. Rapporteur duties were assigned to Dean Courtney, John Hyde, Jae-Bong Lee, Kevin Piner, Darryl Tagami, Kotaro Yokawa, and Lyn Wagatsuma. Wagatsuma served as the lead rapporteur with overall responsibility of assembling the workshop report. Working papers were distributed and numbered (Attachment 2), and the meeting agenda adopted (Attachment 3). All authors who submitted a working paper agreed to have their papers posted on the ISC website where they will be available to the public.

2.0 APRIL 2010 ISC BILLWG WORKSHOP SUMMARY

Gerard DiNardo provided a summary of the intercessional workshop of the ISC BILLWG that was convened in Hakodate, Hokkaido, Japan 15-22 April 2010. The goals of this workshop were to 1) review the status of the North Pacific swordfish assessment using SS3 and Bayesian production models, 2) discuss progress of the blue marlin symposium, 3) delineate striped marlin stock boundaries, and 4) identify potential billfish biological reference points (BRP).

Conclusions from this meeting included:

- The North Pacific striped marlin stock assessment, scheduled to be completed in 2011, would be based on a two stock scenario hypothesis in the North Pacific Ocean. The two stocks are defined by the following boundaries (Figure 1):

- WCPO stock- West of 140°W and north of the equator
 - EPO stock- East of 140°W and north of the equator
- The WG identified 17 potential BRPs for inclusion in the Biological Reference Point Attributes table; these BRPs are commonly used for stock assessment of highly migratory species as discussed during the meeting (Table 2). It was agreed that the each potential BRP should be characterized using the following attributes so that the Northern Committee can understand the implications of each BRP easily: the definition and management purpose, model structure, data needs, limit or target reference point, type of overfishing, pros/cons and special comments (Table 1). It was agreed that the table will be filled out, reviewed and finalized at the July 2010 BILLWG meeting.
 - The North Pacific swordfish Bayesian Surplus Production (BSP) model and Stock Synthesis 3 (SS3) model were updated. Results from the BSP model were similar to the 2009 assessment. Conservation advice will remain unchanged unless clarification is required. The SS3 results in Region 2 do not provide reliable results due to limited data on size at catch.

Figure 1. Stock boundary delineated for the 2011 stock assessment of North Pacific striped marlin.




Table 1. Draft description of biological reference points including definition and management purpose, attributes, and special comments for two example BRPs for exploitation rates.

Biological Reference Point	Definition and Management Purpose	Model Structure	Data Needs	Limit or Target Reference Point	Model Includes Population Dynamics for Recruitment Overfishing	Pros/Cons and Special Comments
FMSY	Fishing mortality that maximizes yield under existing environmental conditions and fishery selectivity pattern	Age-structured or size-structured model for one or two sexes	Fishery catch, fishery catch per unit effort or other relative abundance indices, life history parameters (including natural mortality at age, size at age, weight-length relationships, fishery selectivity pattern, sex ratio in catch if two-sex model)	Has been used as limit and target reference point in various RFMOs	Yes	FMSY is difficult to estimate if stock-recruitment relationship is not known. This BRP may be easy to implement but also entails high risk of recruitment overfishing
FMAX	Fishing mortality that maximizes yield per recruit	Age-structured yield per recruit model	Life history parameters	Has been used as a limit and a target BRP	No	FMAX may be appropriate if recruitment is relatively constant over a range of fishing effort. This BRP may be very risky for some rapidly-growing species because it may cause recruitment overfishing

Table 2. Potential biological reference points for billfish.

Biological Reference Point
F_{MSY}
F_{MAX}
$F_{0.1}$
F_{MED}
F_{τ}
F_{SPR}
$F_{SSB-ATHL}$
F_{lim}
F_{pa}
F_{loss}
B_{MSY}
B_{MAX}
$B_{0.1}$
$B_{X\%}$ (depletion)
B_{lim}
B_{pa}
B_{loss}

2.1 Status of Work Assignments

At the April 2010 workshop, the BILLWG was tasked with a number of assignments that included:

- At the July 2010 BILLWG workshop, Japan, Chinese Taipei, Korea, China, Mexico, USA, and IATTC will present data on the spatial extent and disposition of fisheries catching North Pacific striped marlin within the stock boundaries delineated at April 2010 BILLWG workshop.
- BRP table (Table 1) will be filled out, reviewed and finalized at the next BILLWG meeting in July.
- By the scheduled January 2011 BILLWG workshop, submit stock specific Category I, II, and III North Pacific striped marlin data for review and inclusion in the forthcoming striped marlin stock assessment.

At the April 2010 ISC BILLWG Workshop, the ISC BILLWG Chairman was also tasked with a number of assignments that include:

- Present results from the updated North Pacific swordfish assessment at the 10th ISC Plenary.
- Construct draft outline of proposed objectives and scope for World Blue Marlin Symposium by July 2010 ISC BILLWG workshop.

The WG Chairman reported that not all assignments due at the July 2010 BILLWG workshop were completed. The spatial extent and disposition of fisheries catching striped marlin in the boundary areas was not submitted by all countries. This will impact the North Pacific striped marlin assessment schedule and we look forward to presentations on this topic at the next workshop. The WG Chairman also reminded BILLWG members that some assignments, specifically the submission of stock specific North Pacific striped marlin Category I, II, and III data, are on-going.

Discussion

It was clarified that the stock specific North Pacific striped marlin data should be submitted directly to the BILLWG Chair. The appropriate data will then be passed on to the ISC Database Administrator. It was also clarified that in addition to submitting the Category I, II, and III data, BILLWG members should also submit CPUE time series for use in the stock assessment.

3.0 BIOLOGICAL REFERENCE POINTS

3.1 Age-Based Analyses of Potential Biological Reference Points for the Western and Central North Pacific Swordfish (*Xiphias gladius*) Stock presented by Jon Brodziak (ISC/10/BILLWG-2/02)

Age-based demographic analyses were used to determine a suite of candidate biological reference points for the Western and Central North Pacific swordfish stock for consideration by the ISC Billfish Working Group. Life history data and results from the recent age-structured stock assessment modeling of this stock were used to compute the fishing mortality reference points F_{MSY} , F_{MAX} , $F_{0.1}$, F_{MED} , and F_{SPR} . The same information was used to compute the biomass reference points B_{MSY} , B_{MAX} , $B_{0.1}$, B_{MED} , and B_{SPR} . The percentage of maximum yield and spawning biomass per recruit were summarized to compare the relative yield and stock conservation benefits of the various fishing mortality reference points. Similarly, the ratios of reference biomass, recruitment, and yield to the values at MSY were also summarized to compare the relative stock conservation and yield benefits of the various biomass reference points.

Discussion

It was noted that several choices were made that may affect the resulting BRP estimates. For the example, the choice of years (1994-2006) included in the estimation of average selectivity and the method used to estimate catch weighted quarterly average selectivity. The rationale and/or methodology for these choices should be clarified. It was also noted that the age of the plus group may affect the resulting BRP estimates, especially if fishing mortality is high relative to natural mortality, and that the explicit relationship between the age plus group and BRPs estimated within Stock Synthesis are not well documented. It was noted that the NOAA Fisheries Toolbox (NFT) YPR program was chosen to estimate biological reference points for the working paper because explicit equations for each estimated reference point are available for the NFT YPR program (provided in the working paper).

It was noted that the BRP F_{crash} was in excess of $F=3$ because of the stock recruitment steepness assumption of $h=0.9$.

An observation was made, that based on the current swordfish assessment, the estimate of MSY from SS3 may be too low because fishing pressure is low.

3.2 Production model analyses of maximum sustainable yield-based reference points for the North Pacific swordfish stocks presented by Jon Brodziak (ISC/10/BILLWG-2/03)

Production model analyses of maximum sustainable yield-based reference points were conducted in 2009 and 2010 to assess the Western and Central (WCPO) and the Eastern Pacific (EPO) swordfish stocks in the North Pacific. Estimates of maximum sustainable yield-based reference point from the Bayesian surplus production models of the two swordfish stocks and their variability were summarized for consideration by the ISC BILLWG. The results for the WCPO stock were taken from the 2009 stock assessment. Results for the EPO stock were taken from the 2010 stock assessment update which included an updated time series of swordfish catches in the Eastern Pacific Ocean.

Discussion

It was noted that the estimate of B_{MSY} from the production model differed from the stock synthesis model, and that the difference in B_{MSY} resulted primarily from differences in model structure and the overall lack of contrast in the North Pacific swordfish fisheries CPUE data as discussed at the last working group meeting. In addition, differences in how the yield curve is estimated within age-structured YPR relative to the production model may also affect the resulting estimate of B_{MSY} .

It was noted that within the production model, the estimate of annual harvest rate to produce MSY (H_{MSY}) was higher in the WCPO (0.25) than in the EPO (0.15). There was a discussion about whether or not this difference was real given the uncertainty in H_{MSY} especially within the EPO. Some plausible differences in swordfish habitat between the WCPO and EPO were discussed based on oceanographic differences. It was also discussed that uncertainty about the exact location of the southern boundary for the EPO swordfish stock may also have contributed to the uncertainty of estimated BRPs in the EPO.

3.3 Biological Reference Point Table presented by Kevin Piner and Kotaro Yokawa (ISC/10/BILLWG-2/01)

The completed Biological Reference Point Table that was assigned to be completed at the April 2010 ISC BILLWG workshop was presented for review and finalization. The table includes 17 BRPs that are commonly used for stock assessment of highly migratory species, and were characterized using attributes including: the definition and management purpose, model structure, data needs, limit or target reference point, type of overfishing, pros/cons, and special comments.

Discussion

The BILLWG reviewed the BRP Table and made the following revisions:

- $F_{\infty} = F_{\text{crash}}$
- $F_{\text{SSB}} = F_{\text{SSB-ATHL}}$
- $F_{X\%SPR} = F_{\text{SPR}}$

Merits of each BRP were discussed and included in the table (Table 3).

Table 3. Billfish Biological Reference Point Table

Biological Reference Point	Definition and Management Purpose	Model Structure ¹	Data Needs ²	Limit or Target Reference Point	Type of overfishing	Pros/Cons and Special Comments
F based Reference Points						
F _{MSY}	Fishing mortality that maximizes yield under existing environmental conditions.	Age-structured or size-structured model for one or two sexes	Fishery catch, fishery catch per unit effort or other relative abundance indices, life history parameters	Has been used as limit and target reference point in various RFMOs	Recruitment and growth	F _{MSY} is difficult to estimate if stock-recruitment relationship is not known. This BRP may be easy to implement but also entails high risk of recruitment overfishing. Can be estimated with biomass dynamics modeling.
F _{MAX}	Fishing mortality that maximizes yield per recruit under existing environmental conditions	Age-structured yield per recruit model	Life history parameters	Has been used as a limit and a target BRP	Growth	F _{MAX} may be appropriate if recruitment is relatively constant over a range of fishing effort. This BRP may be very risky for some rapidly-growing species because it may cause recruitment overfishing
F _{0.1}	The fishing mortality rate corresponding to 10% of the slope of the Y/R curve at the origin.	Age-structured yield per recruit model	Life history parameters	Has been used as a limit and a target BRP	Growth	A more precautionary exploitation level relative to F _{MAX} . Often thought to reduce potential recruitment overfishing without a substantial loss in yield.
F _{MED}	The fishing mortality rate to produce replacement recruitment often taken to the median of the R/S distribution. Fishing mortality to maintain recruitment at replacement level observed during specified period.	Estimates of Spawners and Recruits	Estimates of Spawners and Recruits. Typically drawn from an age structured assessment model.	Target or Limit	Recruitment	³ Value dependent on the range of SSB used in the calculations. Not informative if estimates of recruitment taken from a narrow range of spawning biomass. No assumptions about recruitment process. Risky with the specification of BH h=1.0.
F _U	Fishing mortality rate corresponding to the slope of the S/R function at the origin. Theoretical upper	A S/R curve and a relationship of SSB/R	Estimates of Spawner and Recruits. Typically drawn from an age	Limit	Recruitment	Fishing at F _U leads to extinction. Can only be interpreted as a Limit. Upper limit. Does not account for dispensatory effects.

	bound of sustainable rates.	and F	structured assessment model.			
$F_{X\%SPR}$	Fishing mortality rate that produces X% of the unfished spawning potential under equilibrium conditions. Sometimes used for a proxy for other BRP's.	Age-structured Spawner per recruit model	Life history parameters	Has been used as a limit and a target BRP	Recruitment	Although a recruitment based BRP, it is a per-recruit calculation and thus does not depend on estimating the S/R relation. The appropriate level (X%) can be difficult to determine.
F_{SSB}	Fishing mortality rate that produces no more than a specified probability of SSB falling below a defined level of SSB during a given projection period.	Age or length structured assessment	Fishery catch, fishery catch per unit effort or other relative abundance indices. May use additional data such as, life history parameters, biological samples etc.	Target or Limit	Recruitment	Assumes that specified level of spawning biomass is sufficient to insure recruitment success. Flexible which is both a pro and a con. Requires lots of specifications.
F_{lim}	Fishing mortality if maintained will drive the stock to the biomass limit (B_{lim}).	The same as typically associated with an age structured model.	Estimates of spawners and recruits.	Limit	Recruitment	Specified B_{lim} .
F_{pa}	Fishing mortality if maintained drives stock to precautionary biomass limit (B_{pa})	The same as typically associated with an age structured model.	Estimates of spawners and recruits. Need information on accuracy of assessment and risk to be accepted.	Limit	Recruitment	Specified B_{pa} . More precautionary version of F_{lim} .
F_{loss}	Fishing mortality if maintained drives a stock to the lowest observed	Age-structured or size-	Fishery catch, fishery catch per unit effort or other relative	Limit	Recruitment.	Usually used as a proxy of F_{lim} when data is limited.

	spawning stock.	structured model for one or two sexes.	abundance indices, life history parameters.			
Biomass based reference points						
B_{MSY}	The average biomass resulting from fishing at F_{MSY}	Age-structured or size-structured model for one or two sexes.	Fishery catch, fishery catch per unit effort or other relative abundance indices, life history parameters.	Has been used as limit and target reference point in various RFMOs	Recruitment	B_{MSY} is difficult to estimate if stock-recruitment relationship is not known. Can be estimated with biomass dynamics modeling. This BRP may be easy to implement but also entails high risk of recruitment overfishing
B_{MAX}	The average biomass resulting from a fishing mortality that maximizes yield per recruit	Age-structured yield per recruit model	Life history parameters	Has been used as a limit and a target BRP	⁴ Associated value	B_{MAX} may be appropriate if recruitment is relatively constant over a range of fishing effort. Seldom used for management but included because F_{MAX} is defined.
$B_{0.1}$	The average biomass level associated with fishing at $F_{0.1}$	Age-structured or size-structured model for one or two sexes	Fishery catch, fishery catch per unit effort or other relative abundance indices, life history parameters	Has been used as limit and target reference	⁴ Associated value	Seldom used for management but included because $F_{0.1}$ is defined.
$B_{X\%}$ (depletion)	A biomass level that is some specified fraction of the estimated unfished biomass level	Biomass dynamic or age structured model.	Fishery catch, fishery catch per unit effort or other relative abundance indices.	Has been used as limit and target reference	Recruitment	Must use additional analysis to determine the appropriate depletion level. Usually a proxy for B_{MSY} . Depletion is typically calculated relative to unfished level, however substantial uncertainty exists in the calculation of unfished state.
B_{lim}	Set on basis of historical data. Biomass below B_{lim} entails high risk that recruitment might be	The same as typically drawn from an age structured model.	Estimates of spawners and recruits.	Limit	Recruitment	Needs long time series of data (multiple generations). Easy to understand and based on observed values.

	reduced.					
B_{pa}	Precautionary buffer against natural variability and uncertainty associated with B_{lim} . (Note that $B_{pa} > B_{lim}$)	The same as typically drawn from an age structured model.	Estimates of spawners and recruits. Need information on accuracy of assessment and risk to be accepted.	Limit	Recruitment	Needs long time series of data (multiple generations). Easy to understand and based on observed values.
B_{loss}	The lowest observed spawning biomass.	Age structured model.	Fishery catch, fishery catch per unit effort or other relative abundance indices, life history parameters.	Limit	Recruitment.	Used as a proxy for B_{lim} . Needs long time series of data (multiple generations). Easy to understand and based on observed values.

¹Model structure applies to calculation of reference point only. Additional model complexity may be needed to calculate observed metric (F, SSB etc) for comparison.

²Data needs applies to calculation of reference point only. Additional data may be needed to calculate observed metric (F, SSB etc) for comparison.

³ There was no consensus that F_{MED} was risky when steepness was specified as 1.0.

⁴ Associated values are often reported along with their F complement, but may not be used for management.

4.0 BILLFISH CONSERVATION ADVICE

The BILLWG reviewed its previous conservation advice with the objectives of clarifying the statements where necessary. Current conservation advice for the two species follows:

North Pacific striped marlin: “While further guidance from the management authority is necessary, including guidance of reference points and the desirable degree of reduction, the fishing mortality rate of striped marlin (which can be converted into effort or catch in management) should be reduced from the current level (2003 or before), taking into consideration various factors associated with this species and its fishery. Until appropriate measures in this regard are taken, the fishing mortality rate should not be increased”

North Pacific swordfish: “the WCPO and EPO stocks of swordfish are healthy and well above the level required to sustain recent catches”

Discussion

Regarding North Pacific striped marlin, the WG members noted that parts of the statement were ambiguous; however no consensus on clarification of the statement was reached.

Regarding North Pacific swordfish, clarification was sought to define the terms WCPO and EPO, but no consensus was reached. It was noted that these terms were defined in the text describing North Pacific swordfish conservation advice in the previous ISC Plenary report (ISC/09/Plenary/Rep).

5.0 NORTH PACIFIC STRIPED MARLIN REGIONALIZED FISHERIES

5.1 Preliminary analysis of area boundary to standardize CPUE of striped marlin in the North Pacific Ocean presented by Kotaro Yokawa (ISC/10/BILLWG-2/04)

Potential area boundaries in the area west of 140°W of the North Pacific Ocean to standardize CPUE of striped marlin was provided. Spatial patterns were clarified using the delta type, two-step method describing abundance index. Results from the two models used in delta type, two-step method helped to identify optimal area boundaries for latitude and longitude. Summing of AIC for both steps led to the selection of the model in the second step to determine optimal boundaries. It was concluded that choosing the appropriate time series of CPUE is important when using GLM to standardize CPUE or when applying another model like the statHBS. Due to the large effect of gear configuration on CPUE, the method for CPUE standardization of striped marlin should be revisited in the next stock assessment.

Discussion

Since the primary author, Minoru Kanaiwa, was not present to answer questions and to fully explain the model and methodology, it was agreed that full review and discussion on this paper would be postponed until the next ISC BILLWG meeting. Co-author, Kotaro Yokawa, pointed out the difficulties in the estimation of gear configuration within the statistical approach due to

the skewed distribution pattern of data for Japanese offshore and distant-water longliners. This problem should be revisited during the next BILLWG workshop.

5.2 The U.S. Longline Fishery for Striped Marlin in the North Pacific Ocean presented by Gerard DiNardo (ISC/10/BILLWG-2/05)

This report summarizes catch trends for striped marlin caught by the U.S. Hawaii-based longline fishery in the North Pacific Ocean (NPO). Although striped marlin are targeted and taken incidentally by a suite of commercial and recreational fisheries in the North Pacific Ocean, only the U.S. longline fishery is discussed here. To facilitate completion of the upcoming striped marlin stock assessment, which assumes two NPO stocks, the U.S. longline time series for catch has been separated into WCPO and EPO stocks. Trends of catch, number of sets, and number of hooks were presented from 1991-2009. Striped marlin catch was also plotted by area for 2009.

Discussion

It was pointed out that there has been an increasing trend for number of sets and number of hooks since 2001, and that 95% of the effort was in the WCPO. It was also noted that catch varied substantially but was relatively stable from 1991-2009 and nearly all of the catch in 2009 was in the WCPO.

It was suggested that the following summaries be added to the next catch and effort update for striped marlin in the U.S. longline fishery in the NPO:

- Number of vessels by year
- Number of sets and hooks for both shallow and deep sets by year
- Market value by year
- Number of hooks per basket by year

5.3 Available data of striped marlin and swordfish by the Japanese fishery in the North Pacific presented by Kotaro Yokawa (ISC/10/BILLWG-2/06)

This report provides an update of available data for striped marlin by Japanese fisheries, including catch (mt), total hooks, and size data within two-stock structure zones. Catch was estimated separately by gear from Japanese year books and log books, in the WCPO and EPO between 1951 and 2008. Total number of hooks by Japanese offshore longline was estimated in each zone during the same period, as was the number of size samples. Additionally, this study provided the updated catch amount of swordfish in the north Pacific by gear and stock zone. The estimated catch and the total number of hooks of striped marlin in recent years decreased significantly in the two zones, compared to those before 1990. Due to the recent decreasing trend of catch and effort data in the Japanese offshore and distant-water longline fisheries, care should be exercised when using these striped marlin data for stock assessment, especially in the northeastern Pacific Ocean.

Discussion

It was pointed out that the only Japanese fishery in the EPO area, as defined by the BILLWG for the upcoming striped marlin stock assessment, is the distant-water longline fishery. Swordfish catches were also presented by gear from 1951-2008 for both the one stock and two stock scenarios. It was also noted that the number of available size data in the north Pacific decreased substantially since 2004. This may be due to new sampling methods implemented at that time and problems with the choice of fork length measured. Efforts are being made to correct some of these errors which will increase the number of available size data from 2004-2008. The number of sets conducted in the EPO area substantially decreased in recent years. Most of the observed reduction occurred off Mexico, which is the main fishing ground for striped marlin in the EPO. This could have an effect on the representativeness of CPUE obtained from Japanese longline data.

6.0 WORLD BLUE MARLIN SYMPOSIUM

Gerard DiNardo reviewed the rationale, objectives, possible themes, possible sponsors, steering committee members, and timeline of the World Blue Marlin Symposium tentatively scheduled for May 2011.

Discussion

There was discussion on whether holding a World Blue Marlin Symposium (WBMS) is a necessary condition for the completion of a blue marlin stock assessment and it was noted that the assessment would still proceed whether a WBMS was held or not. The BILLWG agreed that although a WBMS would be beneficial, it would not go forward with a WBMS in 2011. It was suggested that input (data) and support from other organizations (i.e. SPC) could be attained by conducting smaller workshops, and not a formal symposium. It was also suggested that the WBMS could be held at a later date, following the completion of a blue marlin stock assessment, possibly affiliated with the International Billfish Symposium.

7.0 OTHER BUSINESS

7.1 RFMO Plans and Upcoming Meetings

7.1.1 IATTC North Pacific striped marlin stock assessment

Gerard DiNardo notified the BILLWG on the IATTC's plan to complete a North Pacific striped marlin stock assessment by the end of July 2010. Further detail is yet unknown. The BILLWG Chair is assigned to look into this, and report back to the BILLWG.

Discussion

Concern was expressed about having two separate North Pacific striped marlin stock assessments. The BILLWG agreed to review the results of the IATTC North Pacific striped marlin stock assessment and decide how to proceed with its stock assessment from there.

7.1.2 International Symposium on Tuna and Billfish Tagging

Gerard DiNardo discussed the upcoming International Symposium on Tuna and Billfish Tagging – Challenges for Tuna and Billfish Tagging Technology and Data Utilization. This symposium is scheduled for 7-12 November 2010 in Taitung, Taiwan. While this is not an “official” ISC symposium, it was noted that some BILLWG members are involved in the planning of this symposium.

Discussion

The BILLWG endorsed the symposium and encourages members to attend. The BILLWG would also like to have a presentation of the results of the symposium at a future BILLWG workshop. Since many billfish stock structure questions still exist, the BILLWG Chairman should consider making a presentation at this symposium on the need for a coordinated billfish tagging program in the Pacific Ocean.

7.2 Work Assignments

The BILLWG were given a number of assignments:

- Submit finalized working papers presented at this meeting by Friday, 13 August 2010.
- All member countries will submit stock specific Category I, II, and III North Pacific striped marlin data as well as CPUE time series for use in the stock assessment by 1 January 2011. All data should be separated into fisheries within the stock boundaries delineated by the BILLWG.
- All member countries will submit Category I data for all billfish (blue marlin, swordfish, etc.) species by 1 January 2011.

The BILLWG Chairman was tasked with a number of assignments:

- Contact WCPFC to request billfish data from non-ISC member countries.
- Determine status of the ISC Biological Sampling proposal and report to BILLWG at January 2011 BILLWG workshop. The Chairman should also seek clarification on status of the Biological Sampling proposal at the 10th ISC Plenary.
- Request information from IATTC regarding North Pacific striped marlin stock assessment and report to BILLWG.

7.3 Future Meetings

The next intercessional BILLWG workshop is scheduled for 19-27 January 2011 in Hawaii, USA. The goals of this workshop will be to review and adopt stock specific Category I, II, and III North Pacific striped marlin data as well as CPUE time series for use in the stock assessment.

All data should be separated into fisheries within the stock boundaries delineated by the BILLWG.

The following intercessional BILLWG workshop is scheduled for 19-27 May 2011. The location is not yet determined, but Japan provisionally offered to host this meeting. The goal of this meeting will be to finalize the North Pacific striped marlin stock assessment for presentation at the 11th ISC Plenary meeting.

8.0 ADJOURNMENT

The ISC BILLWG intercessional workshop was adjourned at 3:37pm on 13 July 2010. The Chairman expressed his appreciation to all participants for their contributions and cooperation in completing a successful meeting.

Table 4. Striped marlin catches (in metric tons) by fisheries, 1952-2005. Blank indicates no effort. - indicates data not available. 0 indicates less than 1 metric ton. Provisional estimates in ().

Year	Japan							Chinese Taipei ¹														Costa Rica ¹	Korea			Mexico			United States					Grand Total										
	Distant-							Coastal Gillnet & Gillnet														Sport	High-seas Drift			Longline	Sport ¹	Total	Longline	Troll	Handline	Sport ¹	Total											
	water and Offshore	Coastal	Other	Small Mesh	Large Mesh			Distant- water	High-seas Drift	Offshore	Offshore	Offshore	Coastal	Coastal	Other	Coastal	Coastal																											
	Longline	Longline	Longline	Gillnet	Gillnet	Other ²	Total	Longline	Gillnet	Longline	Gillnet	Others	Harpoon	Setnet	net	Longline	others	Other	Total				Longline	Gillnet	Total																			
1951	2,494	-	673	-	0	1,281	4,447																											4,447										
1952	2,901	-	722	-	0	1,564	5,187												-															5,210										
1953	2,138	-	47	-	0	954	3,139												-															3,144										
1954	3,068	-	52	-	0	1,088	4,207												-															4,223										
1955	3,082	-	28	-	0	1,038	4,148												-															4,153										
1956	3,729	-	59	-	0	1,996	5,785												-															5,819										
1957	3,189	-	119	-	0	2,459	5,767												-															5,809										
1958	4,106	-	277	-	3	2,914	7,300				543							387	930	-														8,289										
1959	4,152	-	156	-	2	3,191	7,501				391							354	745	-														8,311										
1960	3,862	-	101	-	4	1,937	5,904				398							350	748	-														6,682										
1961	4,420	-	169	-	2	1,797	6,388				306							342	648	-														7,060										
1962	5,739	-	110	-	8	1,912	7,769				332							211	543	-														8,317										
1963	6,135	-	62	-	17	1,910	8,124				560							199	759	-														8,951										
1964	14,304	-	42	-	2	2,344	16,692				392							175	567	-														17,317										
1965	11,602	-	19	0	1	2,794	14,416				355							157	512	-														14,951										
1966	8,419	-	112	0	2	1,570	10,103				370							180	550	-														10,689										
1967	11,698	-	127	0	3	1,551	13,379	2			385							204	591	-														14,019										
1968	15,913	-	230	0	0	1,043	17,186	1			332							208	541	-														17,778										
1969	8,544	600	3	0	3	2,668	11,818	2			571							192	765	-														12,613										
1970	12,996	690	181	0	3	1,032	14,902	0			495							189	684	-														15,604										
1971	10,965	667	259	0	10	2,042	13,943	0			449							135	584	0														14,544										
1972	7,006	837	145	0	243	993	9,224	9			580							126	515	0														9,760										
1973	6,357	632	118	0	3,265	702	11,074	1			368							139	708	0														11,791										
1974	6,700	327	49	0	3,112	775	10,963	24			650							118	792	0														11,810										
1975	5,281	286	38	0	6,534	686	12,825	64			742							96	892	0														13,744										
1976	5,136	244	34	0	3,561	585	9,560	32			347							140	519	0														10,110										
1977	3,019	256	15	0	4,424	547	8,261	17			524							219	760	43														9,105										
1978	3,957	243	27	0	5,593	546	10,366	0			618							78	696	28														11,127										
1979	5,561	366	21	0	2,532	526	9,006	26			432							122	580	-														9,622										
1980	6,378	607	5	0	3,467	536	10,993	61			223							132	416	37														11,479										
1981	4,106	259	12	0	3,866	542	8,785	17			491							95	603	-														9,448										
1982	5,383	270	13	0	2,351	656	8,673	7			397							138	542	39														9,295										
1983	3,722	320	10	22	1,845	827	6,746	0			555							214	769	19														7,573										
1984	3,506	386	9	76	2,257	719	6,953	0			965							330	1,295	23														8,307										
1985	3,897	711	24	40	2,323	733	7,728	0			513							181	694	16														8,498										
1986	6,402	901	33	48	3,536	577	11,497	0			179							148	327	61														11,923										
1987	7,538	1,187	6	32	1,856	513	11,132	31			383							151	565	1														12,029										
1988	6,271	752	7	54	2,157	668	9,909	7			457							169	633	11														11,141										
1989	4,740	1,081	13	102	1,562	537	8,035	8			184							157	349	26														9,098										
1990	2,368	1,125	3	19	1,926	545	5,986	2			137							256	395	315														7,465										
1991	2,845	1,197	3	27	1,302	507	5,881	36			254							286	576	106														7,495										
1992	2,955	1,247	10	35	1,169	303	5,719	1			219							197	417	281														7,400										
1993	3,476	1,723	1	-	828	708	6,736	5			221							142	368	438														8,640										
1994	2,911	1,284	1	-	1,443	383	6,022	1			137							196	334	521														8,479										
1995	3,494	1,840	3	-	970	283	6,590	27			83							82	192	153														8,041										
1996	1,951	1,836	4	-	703	152	4,646	26			162							235	122	429														6,162										
1997	2,120	1,400	3	-	813	163	4,499	59			290							396	138	1,017														6,655										
1998	1,784	1,975	2	-	1,092	304	5,157	90			205							345	144	635														7,053										
1999	1,608	1,551	4	-	1,126	184	4,473	66			128							166	433	433														5,979										
2000	1,152	1,109	8	-	1,062	297	3,628	153			161							369	97	537														5,168										
2001	985																																											

¹ Estimated from catch in

Table 5. Swordfish catches (in metric tons) by fisheries, 1952-2005. Blank indicates no effort. - indicates data not available. 0 indicates less than 1 metric ton. Provisional estimates in ().

Year	Japan								Chinese Taipei ⁵												Korea			Mexico	United States ⁶						Grand Total
	Distant-water and Offshore Coastal Other Bait Fishing Trapnet Other ¹ Total								Distant-water Offshore Offshore Offshore Coastal Coastal Gillnet Coastal Coastal Longline Others Other Total												High-seas Drift			All Gears	Hawaii	California				Total	
																									Longline	Longline	Gill Net	Harpoon	Unknown ⁷		
Year	Longline ²	Longline	Driftnet	Harpoon ³	Fishing	Trapnet	Other ¹	Total	Longline	Longline	Gillnet	Others	Harpoon	Setnet	net	Longline	Others	Other	Total	Longline	Gillnet	Total	All Gears	Longline	Longline	Gill Net	Harpoon	Unknown ⁷	Total		
1951	7,246	115	10	4,131	88	78	10	11,678	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	11,678		
1952	8,890	152	0	2,569	6	68	6	11,691	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	11,691		
1953	10,796	77	0	1,407	20	21	87	12,408	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	12,408		
1954	12,563	96	0	813	104	18	17	13,610	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	13,610		
1955	13,064	29	0	821	119	37	41	14,111	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	14,111		
1956	14,596	10	0	775	66	31	7	15,486	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	15,486		
1957	14,268	37	0	858	59	18	11	15,251	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	15,251		
1958	18,525	42	0	1,069	46	31	21	19,734	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	19,734		
1959	17,236	66	0	891	34	31	10	18,267	-	427	-	-	-	-	91	518	-	-	-	-	-	-	-	-	-	-	-	-	18,785		
1960	20,058	51	1	1,191	23	67	7	21,400	-	520	-	-	-	-	127	647	-	-	-	-	-	-	-	-	-	-	-	-	22,047		
1961	19,715	51	2	1,335	19	15	11	21,147	-	318	-	-	-	-	73	391	-	-	-	-	-	-	-	-	-	-	-	-	21,538		
1962	10,607	78	0	1,371	26	15	18	12,115	-	494	-	-	-	-	62	556	-	-	-	-	-	-	-	-	-	-	-	-	12,671		
1963	10,322	98	0	747	43	17	16	11,244	-	343	-	-	-	-	18	361	-	-	-	-	-	-	-	-	-	-	-	-	11,605		
1964	7,669	91	4	1,006	40	16	26	8,852	-	358	-	-	-	-	10	368	-	-	-	-	-	-	-	-	-	-	-	-	9,220		
1965	8,742	119	0	1,908	26	14	182	10,991	-	331	-	-	-	-	27	358	-	-	-	-	-	-	-	-	-	-	-	-	11,349		
1966	9,866	113	0	1,728	41	11	4	11,763	-	489	-	-	-	-	31	520	-	-	-	-	-	-	-	-	-	-	-	-	12,283		
1967	10,883	184	0	891	33	12	5	12,008	-	646	-	-	-	-	35	681	-	-	-	-	-	-	-	-	-	-	-	-	12,689		
1968	9,810	236	0	1,539	41	14	9	11,649	-	763	-	-	-	-	12	775	-	-	-	-	-	-	-	-	-	-	-	-	12,424		
1969	9,416	296	0	1,557	42	11	14	11,336	0	843	-	-	-	-	7	850	-	-	-	-	-	-	-	-	-	-	-	-	12,186		
1970	7,324	427	0	1,748	36	9	3	9,547	-	904	-	-	-	-	5	909	-	-	-	-	-	-	-	-	-	-	-	-	11,083		
1971	7,037	350	1	473	17	37	31	7,946	-	992	-	-	-	-	3	995	0	-	-	-	-	-	-	-	-	-	-	-	9,044		
1972	6,796	531	55	282	20	1	2	7,687	-	862	-	-	-	-	11	873	0	-	-	-	-	-	-	-	-	-	-	-	8,737		
1973	7,123	414	720	121	27	23	2	8,430	-	860	-	-	-	-	119	979	0	-	-	-	-	-	-	-	-	-	-	-	9,816		
1974	5,983	654	1,304	190	27	16	2	8,176	1	880	-	-	-	-	136	1,017	0	-	-	-	-	-	-	-	-	-	-	-	9,627		
1975	7,031	620	2,672	205	58	18	2	10,606	29	899	-	-	-	-	153	1,081	0	-	-	-	-	-	-	-	-	-	-	-	12,257		
1976	8,054	750	3,488	313	170	14	12	12,801	23	613	-	-	-	-	194	830	0	-	-	-	-	-	-	-	-	-	-	-	13,686		
1977	8,383	880	2,344	201	71	7	2	11,888	36	542	-	-	-	-	141	719	219	-	-	-	-	-	-	-	-	-	-	-	12,961		
1978	8,001	1,031	2,475	130	110	22	1	11,770	-	546	-	-	-	-	12	558	68	-	-	-	-	-	-	-	-	-	-	-	14,049		
1979	8,602	1,038	983	161	45	15	4	10,848	7	661	-	-	-	-	33	701	-	-	-	-	-	-	-	-	-	-	-	-	11,949		
1980	6,005	849	1,746	398	29	15	1	9,043	10	603	-	-	-	-	76	689	64	-	-	-	-	-	-	-	-	-	-	-	10,905		
1981	7,039	727	1,848	129	58	9	3	9,813	2	656	-	-	-	-	25	683	-	-	-	-	-	-	-	-	-	-	-	-	12,820		
1982	6,064	874	1,257	195	58	7	1	8,456	1	855	-	-	-	-	49	905	48	-	-	-	-	-	-	-	-	-	-	-	11,842		
1983	7,692	999	1,033	166	30	9	2	9,931	0	783	-	-	-	-	166	949	11	-	-	-	-	-	-	-	-	-	-	-	12,763		
1984	7,177	1,177	1,053	117	98	13	0	9,635	-	733	-	-	-	-	264	997	48	-	-	-	-	-	-	-	-	-	-	-	13,520		
1985	9,335	999	1,133	191	69	10	0	11,737	-	566	-	-	-	-	259	825	24	-	-	-	-	-	-	-	-	-	-	-	15,981		
1986	8,721	1,037	1,264	123	47	9	0	11,201	-	456	-	-	-	-	211	667	9	-	-	-	-	-	-	-	-	-	-	-	14,761		
1987	9,495	860	1,051	87	45	11	0	11,549	3	1,328	-	-	-	-	190	1,521	44	-	-	-	-	-	-	-	-	-	-	-	15,439		
1988	8,574	678	1,234	173	19	8	0	10,686	-	777	-	-	-	-	263	1,040	27	-	-	-	-	-	-	-	-	-	-	-	14,001		
1989	6,690	752	1,596	362	21	10	0	9,431	50	1,491	-	-	-	-	38	1,579	40	-	-	-	-	-	-	-	-	-	-	-	13,279		
1990	5,833	690	1,074	128	13	4	0	7,742	143	1,309	-	-	-	-	154	1,606	61	-	-	-	-	-	-	-	-	-	-	-	15,672		
1991	4,809	807	498	153	20	5	0	6,292	40	1,390	-	-	-	-	180	1,610	5	-	-	-	-	-	-	-	-	-	-	-	14,306		
1992	7,234	1,181	887	381	16	6	0	9,705	21	1,473	-	-	-	-	243	1,737	8	-	-	-	-	-	-	-	-	-	-	-	19,842		
1993	8,298	1,394	292	309	43	4	1	10,341	54	1,174	-	-	-	-	310	1,538	15	-	-	-	-	-	-	-	-	-	-	-	20,368		
1994	7,366	1,357	421	308	37	4	0	9,493	-	1,155	-	-	-	-	219	1,374	66	-	-	-	-	-	-	-	-	-	-	-	16,228		
1995	6,422	1,387	561	423	34	7	0	8,834	50	1,135	-	-	-	-	225	1,410	10	-	-	-	-	-	-	-	-	-	-	-	14,559		
1996	6,916	1,067	428	597	45	4	0	9,057	9	701	2	-	19	8	-	-	741	15	-	-	-	-	-	-	-	-	-	-	13,957		
1997	7,002	1,214	365	346	62	5	0	8,994	15	1,358	1	1	27	8	-	24	-	100	100	2,365	2,881	512	708	84	11	4	196	17,089			
1998	6,233	1,190	471	476	68	2	0	8,440	20	1,178	8	-	17	15	1	-	-	153	153	3,603	3,263	418	931	48	19	4,679	18,114	18,114			
1999	5,557	1,049	724	416	47	5	0	7,798	70	1,385	4	-	51	5	1	-	-	132	132	1,136	3,100	1,229	606	81	27	5,043	15,625	15,625			
2000	6,180	1,121	808	497	49	5	0	8,660	325	1,531	5	-	74	5	1	1	-	202	2,216	2,949	1,885	646	90	9	5,579	18,929	18,929				
2001	6,932	908	732	230	30	15	0	8,847	1,039	1,691	17	-	64	8	1	1	-	438	438	780	220	1,749	375	52	5	2,401	15,287	15,287			
2002	6,230	965	1,164	201	29	11	0	8,600	1,633	1,557	7	1	1	16	1	1	-	439	439	465	204	1,320	302	90	3	1,919	14,640	14,640			
2003	5,376	1,063	1,198	149	28	4	0	7,818	1,084	2,196	3	-	-	8	-	-	-	381	381	671	147	1,812	216	107	0	2,282	14,443	14,443			
2004	5,395	1,509	1,062	229	30	4	0	8,229	884	1,828	5	-	-	7	1	-	3	410	410	270	213	898	169	62	37	1,379	13,016	13,016			
2005	5,359	1,295	956	187	337	3	0	8,137	437	1,813	1	-	-	5	2	-	18	434	434	235	1,475	-	220	76	0	1,771	12,853	12,853			
2006	6,181	1,508	796	244	342	5	1	9,077	-	-	-	-	-	-	-	-	-	477	477	347	1,175	-	444	71	2	1,692	11,593	11,593			
2007	(6,109)	(2,017)	(829)	(122)	(367)	(2)	(1)	(9,446)	-	-	-	-	-	-	-	-	-	452	452	383	1,444	-	-	484	58	0	1,986	(12,267)	(12,267)		
2008	(4,426)	(1,758)	(648)	(173)	(349)	(3)	(0)	(7,357)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	(7,441)		

¹ Catch data are currently unavailable for Republic of Korea, Philippines, and some other countries catching swordfish in the North Pacific.

² Catches by gear for 1952-1970 were estimated roughly using FAO statistics and other data. Catches for 1971-2002 are more reliably estimated.

³ Constrains trolling and harpoon but majority of catch obtained by harpoon.

⁴ For 1952-1970 "Other" refers to catches by net fishing and various unspecified gears.

⁵ Offshore longline category includes some catches from harpoon and other fisheries but does not include catches unloaded in foreign ports.

⁶ Estimated round weight of retained catch. Does not include discards.

⁷ Unknown includes pole and line, purse seine, troll and troll/handline, half ring, and unspecified gears.

Attachment 1. List of Participants**Japan**

Hideki Nakano
National Research Inst. of Far Seas Fisheries
5-7-1 Orido, Shimizu
Shizuoka, Japan 424-8633
81-54-336-6035, 81-54-335-9642 (fax)
hnakano@affrc.go.jp

Yukio Takeuchi
National Research Inst. of Far Seas Fisheries
5-7-1 Orido, Shimizu
Shizuoka, Japan 424-8633
81-54-336-6039, 81-54-335-9642 (fax)
yukiot@affrc.go.jp

Kotaro Yokawa
National Research Inst. of Far Seas Fisheries
5-7-1 Orido, Shimizu
Shizuoka, Japan 424-8633
81-54-336-6046, 81-54-335-9642 (fax)
yokawa@fra.affrc.go.jp

Republic of Korea

Jae-Bong Lee
National Fisheries Research and Dev. Inst.
152-1, Haean-ro, Gijang-up, Gijang-gun
Busan, Republic of Korea 619-705
82-51-720-2296, 82-51-720-2277 (fax)
leejb@nfrdi.go.kr

Joon-Taek Yoo
National Fisheries Research and Dev. Inst.
152-1, Haean-ro, Gijang-up, Gijang-gun
Busan, Republic of Korea 619-705
82-51-720-2334, 82-51-720-2337 (fax)
yoojt@nfrdi.go.kr

United States

Jon Brodziak
NOAA/NMFS PIFSC
2570 Dole Street
Honolulu, HI 96822-2396
808-983-2964, 808-983-2902 (fax)
Jon.Brodziak@noaa.gov

Dean Courtney
NOAA/NMFS PIFSC
2570 Dole St.
Honolulu, HI 96822-2396
808-983-5345, 808-983-2902 (fax)
Dean.Courtney@noaa.gov

Gerard DiNardo
NOAA/NMFS PIFSC
2570 Dole St.
Honolulu, HI 96822-2396
808-983-5397, 808-983-2902 (fax)
Gerard.DiNardo@noaa.gov

John Hyde
NOAA/NMFS SWFSC
3333 N. Torrey Pines Ct.
La Jolla, CA 92037
858-546-7086, 858-546-7003 (fax)
John.Hyde@noaa.gov

Kevin Piner
NOAA/NMFS PIFSC
2570 Dole St.
Honolulu, HI 96822-2396
808-983-5705, 808-983-2902 (fax)
Kevin.Piner@noaa.gov

Darryl Tagami
NOAA/NMFS PIFSC
2570 Dole St.
Honolulu, HI 96822-2396
808-983-5745, 808-983-2902 (fax)
Darryl.Tagami@noaa.gov

7/14/10

BILLWG

Russ Vetter
NOAA/NMFS SWFSC
8604 La Jolla Shores Drive
La Jolla, CA 92037
858-546-7125, 858-546-5656 (fax)
Russ.Vetter@noaa.gov

Lyn Wagatsuma
Joint Inst. of Mar. and Atmos. Research
2570 Dole St.
Honolulu, HI 96822-2396
808-983-2966, 808-983-2902 (fax)
Lyn.Wagatsuma@noaa.gov

Attachment 2. Working Papers and Background Papers

WORKING PAPERS

ISC/10/BILLWG-2/01	Biological Reference Point Table. Kevin Piner and Kotaro Yokawa. (Kevin.Piner@noaa.gov)
ISC/10/BILLWG-2/02	Age-Based Analyses of Potential Biological Reference Points for the Western and Central North Pacific Swordfish (<i>Xiphias gladius</i>) Stock. Jon Brodziak and Dean Courtney. (Jon.Brodziak@noaa.gov)
ISC/10/BILLWG-2/03	Production model analyses of maximum sustainable yield-based reference points for the North Pacific swordfish stocks. Jon Brodziak and Gakushi Ishimura. (Jon.Brodziak@noaa.gov)
ISC/10/BILLWG-2/04	Preliminary analysis of area boundary to standardize CPUE of striped marlin in North Pacific Ocean. Minoru Kanaiwa and Kotaro Yokawa. (m3kanaiw@bioindustry.nodai.ac.jp)
ISC/10/BILLWG-2/05	The U.S. Longline Fishery for Striped Marlin in the North Pacific Ocean. Russell Ito and Karen Sender. (Russell.Ito@noaa.gov)
ISC/10/BILLWG-2/06	Available data of striped marlin and swordfish by the Japanese fishery in the North Pacific. Ai Kimoto and Kotaro Yokawa. (aikimoto@affrc.go.jp)

BACKGROUND PAPERS

ISC/10/BILLWG-1/REPORT	Report from the April 2010 ISC Billfish Working Group Workshop. 15-22 April 2010. BILLWG. (Gerard.DiNardo@noaa.gov)
------------------------	---

Attachment 3. Agenda

INTERNATIONAL SCIENTIFIC COMMITTEE FOR TUNA AND TUNA-LIKE SPECIES IN THE NORTH PACIFIC

BILLFISH WORKING GROUP (BILLWG)

INTERCESSIONAL WORKSHOP AGENDA

Meeting Site: Hotel Grand Pacific
Galiano Room #233
463 Belleville Street
Victoria BC Canada, V8V 1X3
250-386-0450, 250-380-4475 (fax)

Meeting Dates: 12-13 July 2010

Goals: Finalize advice on potential billfish biological reference points, delineate spatial structure of North Pacific striped marlin within areas (WCPO and EPO) delineated at the April 2010 BILLWG Workshop, review and modify (if necessary) current conservation advice, and review World Blue Marlin Symposium proposal.

July 12 (Monday), 0830-0900 – Registration

July 12 (Monday), 0900-1200

1. Opening of Billfish Working Group (BILLWG) Workshop
 - a. Welcoming Remarks
 - b. Introductions
2. Adoption of Agenda and Assignment of Rapporteurs
3. Computing Facilities
 - a. Access
 - b. Security Issues
4. Numbering Working Papers and Distribution Potential
5. Status of Work Assignments and Meeting Summaries
6. Biological Reference Points
 - a. Potential BRP for swordfish
 - b. Finalize BRP table

July 12 (Monday), 1200-1300 – Lunch

July 12 (Monday), 1300-1400

7. Review of Current Billfish Conservation Advice

July 12 (Monday), 1400-1700

8. Description of Regionalized Fisheries
 - a. Country reports

July 13 (Tuesday), 1000-1100

9. World Blue Marlin Symposium Planning
10. Other Matters
 - a. RFMO Plans
 - b. Work Assignments
 - c. Future Meetings

July 13 (Tuesday), 1100-1400

11. Rapporteurs complete sections & report finalized

July 13 (Tuesday), 1400-1600

12. Clearing of Report
13. Adjournment