



Japanese troll CPUE targeting age-0 Pacific bluefin tuna: Updated up to 2014 fishing year

Osamu Sakai and Kazuhiro Oshima

National Research Institute of Far Seas Fisheries,
Fisheries Research Agency
5-7-1, Orido, Shimizu-ku, Shizuoka 424-8633, JAPAN

November 2015

Information paper submitted to the ISC Pacific bluefin tuna Working Group, International Scientific Committee for Tuna and Tuna-Like Species in the North Pacific Ocean (ISC), 18-25 November 2015, Kaohsiung, Taiwan.
Document not to be cited without author's permission.

Summary

The estimate of relative recruitment abundance index from troll fishery in the East China Sea (coastal area around Nagasaki prefecture) in 2014 fishing year was a record-low. This recruitment index was standardized CPUE for troll fishery during 1980-2014 fishing year. Generalized liner model (GLM) with lognormal error distribution was applied for standardization which was authorized and used in previous studies (Ichinokawa et al. 2012, Oshima et al. 2013, Fujioka et al., 2014). This index is a candidate of the abundance index as an input data for upcoming stock assessment of Pacific bluefin tuna.

Introduction

The index of juvenile Pacific bluefin tuna (PBF) abundance based on catch and effort data of a troll fishery is one of the important indices available for monitoring and assessment of the PBF stock. Ichinokawa et al. (2012) provided three CPUE series of troll fisheries from Kochi, Wakayama, and Nagasaki Prefectures, and ISC PBFWG decided to fit only Nagasaki's series in the assessment model due to representativeness (ISC 2014). This troll fishery is targeting age-0 PBF which comes from both two major spawning grounds (in waters near the Ryukyu Islands to the east of Taiwan, and in the southern portion of the Sea of Japan), thus their CPUE would reflect the whole annual recruitment strength of PBF population.

This document presents an update of the standardized CPUE of troll fishery of Nagasaki Prefecture for the upcoming stock assessment scheduled in next February. The catch-and-effort data used in this document have been collected and archived by National Research Institute of Far Seas Fisheries with cooperation from local fishery institutes, as a part of the Marine Ranching Project during 1980's (Secretariat of Forestry and Fisheries Research Council 1989) and Research Project on Japanese bluefin tuna (RJB) since 1994 (Ichinokawa et al. 2012).

Methods

Troll CPUE was based on the catch-and-effort data which have been collected at the 5 main fishing ports in Nagasaki Prefecture since 1980s; i.e. Izuhara-Are, Kami-tsushima, Kami-agata, Ojikam and Tomie. These ports are located in Goto islands and Tsumi islands of western part of Japan (Fig. 1). The unit of the catch and effort is landing weight and number of ship-landing per day, respectively.

Generalized liner model (GLM) with lognormal error distribution was applied to standardize the CPUE, because the effort data (ship-landing information) was not recorded for zero-catch trip. The following three effects were used for the standardization;

- 1) FISHING YEAR (FY); 1980-2014... Fishing year is starting July and ending to June.
- 2) FISHING MONTH (FM); 1-12... Fishing months are aligned with fishing year.
- 3) PORT; five ports... Izuhara-Are, Kami-tsushima, Kami-agata, Ojika, and Tomie.

Objective variable was log(CPUE) and explanatory variables were the three effects listed above and all possible first-order interactions. The GLM was carried out through GLM procedure of SAS 9.4. The best model was determined based on BIC. The standardized CPUE was calculated from

least square mean of 'FY' effect.

Results and Discussions

Catch-and-effort data by each landing port are summarized in Table 1. In 2014 fishing year, total catch amounts and efforts were substantially smaller than previous years in all five ports. This suggests that the fishermen didn't go PBF fishing very much, and/or there were a lot of unrecorded zero-catch trips during 2014 fishing year.

The best model selected by BIC was a combination of "FY", "FM" and "PORT" (Table 2). Residuals distributed centrally around zero, although those distributions showed slight left-skewed shapes (Figs 2 and 3). Range of CV of standardized CPUE was 0.02-0.05. These results and past year trend were not different from the previous study (Fujioka et al. 2014). Updated data points of both nominal and standardized CPUEs showed the increase of recruitment from 2012 to 2013, but decreased again in 2014 fishing year (Fig 4). The 2014 values for both nominal and standardized CPUE were the lowest ever recorded: at less than half of the historical average level.

Those results suggest that intensity of 2014 year class was substantially lower level. This supports the results of "quick estimations of recruitment abundance" which was presented by Japan during April WG (ISC 2015). Thus the "quick estimations" would be of considerable importance as the "warning sign" to detect the most recent recruitment level precociously.

This updated troll index doesn't reflect the recruitment status in most recent fishing year (2015 fishing year: from July 2015). Given the importance of recruitment in influencing stock biomass, the "quick estimations of recruitment abundance" should be noted continually to monitor and consider most recent recruitment level.

References

- Fujioka, K., Oshima, K., Tei, Y., Ichinokawa, M., and Takeuchi Y. 2014. Recruitment abundance indices of young Pacific bluefin tuna revealed by Japanese Troll Fisheries. ISC/14/PBFWG-1/07.
- Ichinokawa, M., Oshima, K. and Takeuchi Y. 2012. Abundance indices of young Pacific bluefin tuna, derived from catch-and-effort data of troll fisheries in various regions of Japan. ISC/12-1/PBFWG/11.
- ISC 2014. Stock assessment of bluefin tuna in the Pacific Ocean in 2014. Report of the Pacific bluefin tuna working group. 121p.
- ISC 2015. Report of the Pacific bluefin tuna working group intersessional workshop. Annex 9 of Report of the fifteenth meeting of the international scientific committee for tuna and tuna-like species in the North Pacific Ocean, plenary session. 24p.
- Oshima, K., Fujioka, K., Ichinokawa, M., and Takeuchi Y. 2013. Updated Japanese troll CPUE targeting age 0 PBF through 2011. ISC/13/PBFWG/Appendix C.

Table 1 Total catch (mt), effort (number of landing per day, excluding zero PBF catch) and CPUE (kg/landing) by year and by fishing port, recorded in catch-and-effort data used for standardization of CPUE in Nagasaki Prefecture.

Fishing year	Catch (mt)						Effort (Number of landing)						CPUE (kg / landing)					
	Tomie	Are	Kami-tsushima	Kami-agata	Ojika	Total	Tomie	Are	Kami-tsushima	Kami-agata	Ojika	Total	Tomie	Are	Kami-tsushima	Kami-agata	Ojika	Total
1980	210.4	7.2	11.2	18.2	11.4	258.3	5330	670	142	339	723	7204	39.5	10.7	78.7	53.7	15.7	35.9
1981	423.0			118.1	125.7	666.8	9740			1633	2952	14325	43.4			72.3	42.6	46.5
1982	62.5	14.3	8.9	45.9	17.9	149.5	1301	694	274	1503	725	4497	48.1	20.7	32.4	30.5	24.7	33.3
1983	242.9	51.3	153.4	350.9	102.4	900.9	6264	1756	2012	3958	2278	16268	38.8	29.2	76.2	88.7	45.0	55.4
1984	482.2	72.8	63.5	355.0	132.6	1106.2	12383	1591	1130	6715	3381	25200	38.9	45.8	56.2	52.9	39.2	43.9
1985	182.7	78.3	85.0	130.8	91.4	568.1	6932	1753	1035	2470	1787	13977	26.4	44.6	82.1	53.0	51.1	40.6
1986	378.5	67.0	24.0	130.5	77.3	677.2	11457	1729	338	2420	2367	18311	33.0	38.7	70.9	53.9	32.6	37.0
1987	115.1	14.3	23.2	132.3	15.1	300.0	4406	500	447	2502	658	8513	26.1	28.6	51.8	52.9	23.0	35.2
1988	281.2	6.0	37.3	150.3	51.1	525.9	9115	283	555	2465	1079	13497	30.8	21.1	67.3	61.0	47.3	39.0
1989	119.5	17.4	36.1	81.2	24.8	279.0	5744	776	696	1583	868	9667	20.8	22.4	51.8	51.3	28.6	28.9
1990	240.9	46.3	145.4	173.2		605.8	6733	903	1537	1739		10912	35.8	51.3	94.6	99.6		55.5
1991	79.0	44.0	95.5	111.7	127.1	457.3	1546	865	1008	1603	2195	7217	51.1	50.9	94.7	69.7	57.9	63.4
1992	66.4	1.9	23.1	12.9	15.1	119.4	2416	234	630	446	953	4679	27.5	8.0	36.7	29.0	15.9	25.5
1993	42.4	17.8		60.1	4.9	125.2	1810	986		2040	487	5323	23.4	18.0		29.4	10.1	23.5
1994	464.1	105.3		874.2	426.3	1869.9	5363	1343		5719	3668	16093	86.5	78.4		152.9	116.2	116.2
1995	104.6			243.4	41.0	389.0	2981			2055	1116	6152	35.1			118.4	36.7	63.2
1996	340.5	104.5		507.1	127.6	1079.8	6134	1543		4793	2065	14535	55.5	67.7		105.8	61.8	74.3
1997	90.4	23.4	59.1	138.8	39.5	351.2	2334	761	690	2605	767	7157	38.7	30.7	85.6	53.3	51.6	49.1
1998	234.3	45.4	196.0	268.8	21.5	766.0	4525	1236	2348	3908	399	12416	51.8	36.7	83.5	68.8	53.9	61.7
1999	202.0	101.8		355.9	74.7	734.4	4294	1167		2691	833	8985	47.1	87.3		132.3	89.6	81.7
2000	48.4	113.4	207.2	318.3	48.2	735.5	2571	1213	1353	2216	668	8021	18.8	93.5	153.1	143.6	72.2	91.7
2001	87.5	76.4	163.8	159.3	48.0	535.1	1582	1111	1682	1729	776	6880	55.3	68.8	97.4	92.2	61.8	77.8
2002	105.5	34.5	44.4	69.1	24.6	278.2	2725	902	951	1495	806	6879	38.7	38.3	46.7	46.2	30.5	40.4
2003	18.0	30.0	68.5	8.1	13.0	137.6	853	631	842	239	357	2922	21.2	47.6	81.3	33.8	36.4	47.1
2004	117.5	83.4	188.2	324.1	40.0	753.3	2304	923	1478	3101	692	8498	51.0	90.4	127.3	104.5	57.7	88.6
2005	22.5	15.2	125.9	68.2	23.6	255.4	550	365	1014	721	354	3004	40.9	41.8	124.1	94.6	66.5	85.0
2006		9.5	30.7	20.0	0.4	60.7		231	437	490	28	1186		41.3	70.4	40.8	16.0	51.2
2007	5.3	22.6	91.8	163.8	29.8	313.3	64	376	753	1920	393	3506	82.6	60.1	121.9	85.3	75.9	89.4
2008	179.7		142.0	53.8	60.9	436.3	2668		854	760	792	5074	67.3		166.3	70.8	76.9	86.0
2009	97.3	35.7	75.6		5.3	213.9	1339	743	693		175	2950	72.7	48.0	109.1		30.1	72.5
2010	115.3	14.7	76.7	171.9	6.5	385.1	2119	439	806	2350	135	5849	54.4	33.4	95.1	73.2	48.4	65.8
2011	28.7	13.2	96.9	216.6	1.7	357.0	979	195	665	2286	55	4180	29.4	67.4	145.7	94.7	31.2	85.4
2012	7.7	18.9	0.6	62.2	3.2	92.6	234	764	19	1526	94	2637	32.9	24.7	33.6	40.8	33.6	35.1
2013	98.0	63.3	6.8	179.7	12.0	359.8	1240	1033	144	1742	179	4338	79.0	61.3	47.1	103.1	67.1	82.9
2014	8.2	3.7	0.0	0.1	0.7	12.7	312	193	4	5	37	551	26.4	19.1	11.5	16.7	18.7	23.1

Table 2 Values of BIC (Bayesian Information Criterion) calculated for all models of possible combinations of main effects and first-order interaction terms. The first best model “a)” (shaded) is used for base case.

Model	BIC
a) fy+fm+port	<u>29157.9</u>
b) fy*fm+port	29186.9
c) fy*port+fm	29412.3
d) fy+fm*port	29202.7
e) fy*fm+fy*port	29629.9
f) fy*fm+fm*port	29204.9
g) fy*port+fm*port	29413.3
h) fy*fm+fm*port+fy*port	29533.6

Table 3 Type 3 analysis of the explanatory variables in the final model for CPUE standardization.

Effects	df	Type III SS	Mean square	F value	Pr > F
Model	48	2520.8	52.5	54.66	<.0001
Error	10225	9823.9	1.0		
Corrected Total	10273	12344.7			
				R squared value	0.204

Effects	df	Type III SS	Mean square	F value	Pr > F
fy	34	1288.0	37.9	39.4	<.0001
fm	10	244.5	24.5	25.5	<.0001
port	4	859.0	214.8	223.5	<.0001

Table 4 Standardized troll CPUE in Nagasaki Prefecture, comparing with previous result (Fujioka et al. 2014) and nominal CPUE. All CPUEs are normalized by each average.

Fishing year	Nominal CPUE	Record Number	Standardized CPUE				Fujioka et al. (2014)	
			Estimation	CV	Lower 5%	Upper 5%	Estimation	CV
1980	0.59	255	0.66	0.03	0.55	0.80	0.65	0.04
1981	0.90	265	1.16	0.03	0.97	1.40	1.14	0.03
1982	0.57	183	0.61	0.04	0.50	0.75	0.60	0.04
1983	0.89	328	0.91	0.03	0.77	1.09	0.89	0.03
1984	0.75	396	0.93	0.03	0.79	1.10	0.91	0.03
1985	0.83	375	0.87	0.03	0.73	1.03	0.85	0.03
1986	0.72	492	0.98	0.02	0.83	1.16	0.96	0.03
1987	0.60	310	0.71	0.03	0.59	0.85	0.70	0.04
1988	0.71	356	0.82	0.03	0.69	0.97	0.80	0.03
1989	0.52	351	0.65	0.03	0.55	0.77	0.64	0.04
1990	1.18	333	1.28	0.02	1.08	1.53	1.25	0.03
1991	1.11	271	1.31	0.03	1.09	1.58	1.28	0.03
1992	0.49	308	0.58	0.03	0.48	0.69	0.56	0.04
1993	0.41	330	0.49	0.03	0.41	0.58	0.48	0.04
1994	1.80	439	2.02	0.02	1.71	2.38	1.98	0.03
1995	0.96	243	1.08	0.03	0.90	1.31	1.06	0.03
1996	1.21	448	1.61	0.02	1.36	1.90	1.58	0.03
1997	0.95	251	0.95	0.03	0.79	1.14	0.92	0.03
1998	0.98	350	0.83	0.03	0.70	0.99	0.81	0.03
1999	1.49	286	1.52	0.02	1.27	1.82	1.49	0.03
2000	1.57	273	1.16	0.03	0.97	1.39	1.14	0.03
2001	1.31	265	1.16	0.03	0.97	1.40	1.14	0.03
2002	0.73	275	0.75	0.03	0.63	0.90	0.74	0.04
2003	0.81	184	0.65	0.03	0.53	0.79	0.64	0.04
2004	1.51	369	1.29	0.02	1.09	1.54	1.27	0.03
2005	1.69	230	1.42	0.03	1.18	1.72	1.39	0.03
2006	1.06	106	0.73	0.04	0.58	0.93	0.71	0.04
2007	1.60	244	1.41	0.03	1.17	1.70	1.37	0.03
2008	1.68	285	1.43	0.02	1.19	1.71	1.40	0.03
2009	1.01	206	1.16	0.03	0.95	1.40	1.13	0.03
2010	1.10	324	1.12	0.03	0.94	1.33	1.09	0.03
2011	1.39	266	0.96	0.03	0.80	1.16	0.94	0.03
2012	0.49	242	0.50	0.03	0.42	0.60	0.51	0.04
2013	1.07	338	0.84	0.03	0.71	0.99		
2014	0.34	97	0.42	0.05	0.33	0.53		




Fig. 1 Location of fishing ports where catch-and-effort data of troll fisheries have been collected in Nagasaki Prefecture.




Fig. 2 Standardized residuals (left panel) and Q-Q plot of them (right panel).




Fig. 3 Standardized residuals by year.




Fig. 4 Comparison of time series of CPUE. Gray and black lines indicate nominal and standardized CPUE from 1980 to 2014 fishing year. Solid circles show the previous standardized CPUE estimated by Fujioka et al. (2014).